

Original Article

Effect of quadriceps and calf muscles fatigue on standing balance in healthy young adult males

A.A. Joudeh¹, A.H. Alghadir¹, H. Zafar¹, S.M. Elwatidy², C. Tse¹, S. Anwer¹

¹Department of Rehabilitation Sciences, College of Applied Medical Sciences, King Saud University, Riyadh, Saudi Arabia; ²College of Medicine, King Saud University, Riyadh, Saudi Arabia

Abstract

Objective: The present study aimed to compare the effects of quadriceps or calf muscles fatigue on static and dynamic standing balance in young healthy adult males. **Methods:** Forty-five healthy, physically active male adults aged 18-30 years were randomly divided into three groups; Quadriceps muscle fatigue group ($n=15$), Calf muscles fatigue group ($n=15$), and a control group ($n=15$). The Modified Clinical Test of Sensory Interaction on Balance, Unilateral Stance, and Limits of Stability (reaction time and movement velocity) were selected as outcome measures for this study. **Results:** The results showed a non-significant difference between pretest and posttest balance scores ($p>0.05$) for quadriceps and calf muscles fatigue on balance. Similarly, there were non-significant differences in posttest balance scores when comparing fatigue effects between the groups ($p>0.05$). **Conclusions:** These results suggested that the fatigue of the quadriceps or calf muscles did not influence standing balance in healthy young adult males. Future longitudinal studies are recommended to further understanding the mechanisms behind localized muscle fatigue effects on standing balance in subjects of different age groups of both genders.

Keywords: Fatigue, Balance, Limit of Stability, Male

Introduction

Decreased balance is common among the elderly people whose postural control system exhibits reduced sensitivity due to deficits in their sensory systems^{1,2}. There are various factors that affect balance such as proprioceptive deficits and muscle weakness³. When considering the relationship between muscle weakness and balance, the weakness of the hamstring, calf, and quadriceps muscles are important⁴. This can have an adverse effect, as maximal muscle strength of the lower limbs is necessary for maintaining optimal postural control, and poor strength levels have been commonly accepted as risk factors which can contribute to falls in the elderly⁵.

Muscle fatigue is another factor that may affect postural control in healthy young and elderly population resulting increased risk of fall⁶⁻⁸. In addition, muscle fatigue can further increase the risk of injury⁹. Muscle fatigue is characterized as diminished ability of the muscle to contract and apply a force that develops immediately after the onset of the sustained physical activity^{10,11}. One of the outcomes of fatigue is that it minimizes the skill-related physical fitness performance, for example, balance¹², strength^{9,12}, and agility¹³ which thusly affect sports performance. A study showed that the lower-limb muscular fatigue caused increased postural sway in

The authors have no conflict of interest.

Corresponding author: Shahnawaz Anwer, MPT, Researcher, Department of Rehabilitation Sciences, CAMS, King Saud University, Riyadh, Saudi Arabia
E-mail: anwer_shahnawazphysio@rediffmail.com

Edited by: F. Rauch
Accepted 8 August 2017

Abbreviations

| |
|--|
| BMI = body mass index |
| mCTIB = Modified Clinical Test of Sensory Interaction on Balance |
| US = Unilateral Stance |
| EO = Eyes open |
| EC = Eyes closed |
| LOS = Limits of Stability |
| RT = Reaction time |
| MVL = Movement velocity |

young people¹⁴. Moreover, fatigue of postural back muscles caused decreased head and trunk balance control and the quadriceps muscular fatigue affects the gait parameters related to slip propensity¹⁵. However, in both young and older people, the effects of fatigue of specific muscle groups on muscular strength have been widely investigated¹⁶. In this respect, despite age-related loss or reduction in muscle strength, previous studies have demonstrated that elderly people show better muscle fatigue resistance than young people following dynamic and isometric lower extremity muscle tests¹⁷⁻¹⁹.

It is not clear how localized muscle fatigue may affect the balance. Moreover, due to muscular fatigue, increased joint laxity indirectly leads to alterations in joint kinesthesia and position sense²⁰. It is suggested that muscle fatigue may decrease muscle response to neural excitation²¹ and decreased contraction activities of the muscles²². In this respect, it can disrupt afferent feedback and alters joint awareness. However, fatigue may occur as a result of an interruption to the chain of events from the central nervous system to the muscle fiber. Previous studies have reported that muscle fatigue negatively affects the balance and control of body position^{17,20}. Researchers have investigated ways in which fatigue affects balance by applying strenuous aerobic exercise or selective muscle-fatiguing protocols to induce generalized muscle fatigue^{9,11}.

Centil et al.¹⁷ investigated the fatigue effect of lower-extremity and trunk muscles on the balance in healthy adults. They reported an impaired balance following a generalized fatigue protocol of the lower-extremity and trunk muscles. Another study additionally proposed that muscle fatigue appears to affect active joint reposition sense²³. Accordingly, muscle fatigue causes muscle spindle information less effective. More recently, Nam et al.⁷ investigated the fatigue effect of gastrocnemius muscle on postural control. They reported impaired medio-lateral stability following gastrocnemius muscle fatigue in the elderly. The detrimental effects of localized muscle fatigue on static balance have been investigated^{4,24}, but its effects on measures of dynamic balance are unknown. This study aims to compare the effects of quadriceps or calf muscles fatigue on static and dynamic standing balance in healthy young adult males.

Materials and methods

Study design

This was a randomized controlled study. The independent variables were (a) fatigue, which included the quadriceps muscle fatigue and calf muscles fatigue and, (b) non-fatigue. Balance indices (Modified Clinical Test of Sensory Interaction on Balance, Unilateral Stance, and Limits of Stability) measured using the Balance Master were the dependent variables.

Participants

Healthy male adults aged 20-30 years from the 'College of Applied Medical Sciences' at 'King Saud University', Riyadh,

participated in this study. Subjects were excluded if they had any history of musculoskeletal, neurological, auditory and visual, cardiovascular, respiratory disorders, and were current smokers, or obese individuals [body mass index (BMI)>35 kg/m²].

Sample size calculation

Based on the results of power analysis, a total sample of forty-five subjects was required to achieve 80% power at a p value of less than 0.05. The sample size was calculated using GraphPad StatMate version 2.00 for Windows (GraphPad Software, San Diego California USA).

Procedure

After a brief description of the study, subjects who fulfilled the inclusion criteria were asked to provide written informed consent. Subjects had the ability to withdraw from the study at any time. The study was approved by the ethics committee of the College of Applied Medical Sciences, King Saud University. After initial evaluation and pretest, participants were allocated randomly to group A (quadriceps muscle fatigue), group B (calf muscles fatigue) or group C (non-fatigue) by choosing an opaque envelope, which contained the names of one of the three groups.

The participants in the fatigue groups performed 5 minutes of warming up on a stationary bicycle at a self-selected speed. One maximal repetition for quadriceps muscle was measured on quadriceps exercise bench²⁵. The fatiguing exercises took place beside the Balance Master in order to obtain a very short lag time between the exercise induced fatiguing activity and the Balance Master assessment²⁶. The average approximate time between the fatiguing process and measurement was 20 seconds. The rating of perceived exertion from the participants was assessed prior to the fatigue intervention to indicate the level of fatigue²⁷. A second rating of perceived exertion was recorded as soon as the participants completed the fatigue protocol. Adequate fatigue was deemed to be a rating of perceived exertion of 14 to 17, which represents 75% to 90% maximum oxygen consumption posttest^{27,28}.

In order to induce fatigue in the quadriceps muscles, the participants were asked to perform knee extensions on the quadriceps exercise bench. They performed two sets of 50 repetitions each with 50% of one maximal repetition and a 4 minute rest interval between the sets²⁵. When the participants could not continue the 50 repetitions in the first set, the fatigue protocol was stopped and the balance test administered. However, if they could complete the first set of 50 repetitions, they continued the protocol after a 4-minute rest²⁵.

Calf muscle fatigue was achieved by asking the subject to perform toe lifts carrying sand weights (10% of body weight) on his shoulders. The fatigue level was reached when subjects were no longer able to perform or finish two sets of 50 repetitions with 4-minute rest between sets^{29,30}. This protocol was chosen based on previous studies with

Table 1. Effect of quadriceps muscle fatigue on balance scores.

| Variables | Pretest Mean ± SD | Posttest Mean ± SD | T test P |
|-----------------------------|----------------------|-----------------------|-------------|
| mCTIB Firm - EO (deg/sec) | 0.42 ± 0.50 | 0.45 ± 0.15 | 0.788 |
| mCTIB Firm - EC (deg/sec) | 0.31 ± 0.08 | 0.35 ± 0.09 | 0.164 |
| mCTIB Foam - EO (deg/sec) | 0.77 ± 0.22 | 0.77 ± 0.20 | 1 |
| mCTIB Foam - EC (deg/sec) | 1.43 ± 0.43 | 1.33 ± 0.39 | 0.29 |
| US - EO Left (deg/sec) | 0.96 ± 0.22 | 1.16 ± 0.58 | 0.154 |
| US - EC Left (deg/sec) | 2.87 ± 0.89 | 2.80 ± 0.82 | 0.63 |
| US - EO Right (deg/sec) | 0.97 ± 0.28 | 1.17 ± 0.39 | 0.101 |
| US - EC Right (deg/sec) | 2.76 ± 1.13 | 3.06 ± 0.93 | 0.285 |
| LOS - RT Forward (sec) | 1.12 ± 0.34 | 0.88 ± 0.34 | 0.223 |
| LOS - RT Right (sec) | 0.90 ± 0.28 | 0.85 ± 0.31 | 0.453 |
| LOS - RT Back (sec) | 0.79 ± 0.27 | 0.78 ± 0.32 | 0.859 |
| LOS - RT Left (sec) | 0.94 ± 0.30 | 0.92 ± 0.23 | 0.833 |
| LOS - MVL Forward (deg/sec) | 3.3 ± 0.91 | 3.44 ± 1.61 | 0.69 |
| LOS - MVL Right (deg/sec) | 4.11 ± 1.28 | 4.60 ± 1.48 | 0.139 |
| LOS - MVL Back (deg/sec) | 3.02 ± 1.08 | 2.89 ± 0.83 | 0.641 |
| LOS - MVL Left (deg/sec) | 4.8 ± 1.58 | 5.06 ± 1.72 | 0.305 |

mCTIB, Modified Clinical Test of Sensory Interaction on Balance; US, Unilateral Stance; EO, Eyes open; EC, Eyes closed; LOS, Limits of Stability; RT, Reaction time; MVL, Movement velocity.

few modifications to induce similar types of fatigue in both muscle groups.

For the non-fatigue group, the participants sat on a chair for 15 minutes between the pretest and posttest balance assessments.

Outcome assessment

The 'Modified Clinical Test of Sensory Interaction on Balance' (mCTSIB) is a modified version of the original Clinical Test of Sensory Interaction on Balance³¹. It was designed to provide information about a patient's functional balance control and to assess postural stability during four sensory conditions including: (1) eyes open on a firm surface, (2) eyes closed on a firm surface, (3) eyes open on foam, and (4) eyes closed on the foam. The mCTSIB test provides an objective measure of patient sway velocity (degrees per second) for each of the aforementioned four task conditions³¹. The Unilateral Stance (US) is a test to investigate the postural stability during one leg stance with the eyes open and closed (ICC=0.79 to 0.95)³². The US improves on the observational testing of unilateral standing performance by providing an objective measure of sway velocity (in degrees per second). Subjects performed three trials for each of the task conditions³³. The Limits of Stability (LOS) is a performance assessment based on a set of motor tasks that identify the total distance the patient can purposely displace his center of gravity without losing balance (ICC=0.82-0.48)¹⁴. It has the capacity to assess the patient's ability to perform specific dynamic balance tasks while monitoring a real-time display of the body's center of gravity position with regards to targets centered over a given base of support and

at the stability limits. The test objectively measures movement reaction time (sec) and movement velocity (degrees per second) for each direction³⁴. All the tests were measured at baseline and posttest after the fatigue protocol.

Statistical analysis

All the data were analyzed using SPSS software version 19 (SPSS, Chicago, Illinois). Results were reported as mean ± standard deviation. Data were tested using the paired t-tests to compare between and within group differences for balance indices ('modified clinical test of sensory interaction on balance', 'unilateral stance', and 'limits of stability'), respectively. The probability level for all the tests was set at 0.05 to indicate significance.

Results

The demographic parameters of the participants between the Quadriceps Fatigue (age, 21.4±2.2 years; height, 1.71±0.06 m; weight, 72.4±13.1 kg; BMI, 25.2±4.6 kg/m²), Calf Fatigue (age, 21.6±2.1 years; height, 1.70±0.04 m; weight, 69.7±14.8 kg; BMI, 24.2±4.7 kg/m²), and Non-Fatigue groups (age, 21.2±0.8; height, 1.71±0.01; weight, 75.3±19.6; BMI, 25.5±6.2) showed no significant difference between the groups.

For the Quadriceps Fatigue Group, the effect of fatigue on balance scores (mCTSIB, US, and LOS) showed no significant difference in all tested positions ($p>0.05$) (Table 1). Similarly, in the Calf Fatigue Group, the effect of

Table 2. Effect of calf muscle fatigue on balance scores.

| Variables | Pretest Mean ± SD | Posttest Mean ± SD | T test |
|-----------------------------|----------------------|-----------------------|--------|
| | | | P |
| mCTIB Firm - EO (deg/sec) | 0.39 ± 0.20 | 0.44 ± 0.12 | 0.396 |
| mCTIB Firm - EC (deg/sec) | 0.31 ± 0.05 | 0.5 ± 0.50 | 0.162 |
| mCTIB Foam - EO (deg/sec) | 0.68 ± 0.12 | 0.68 ± 0.10 | 1 |
| mCTIB Foam - EC (deg/sec) | 1.41 ± 0.27 | 1.25 ± 0.36 | 0.112 |
| US - EO Left (deg/sec) | 0.84 ± 0.17 | 0.81 ± 0.08 | 0.43 |
| US - EC Left (deg/sec) | 2.37 ± 0.62 | 2.13 ± 0.90 | 0.395 |
| US - EO Right (deg/sec) | 0.82 ± 0.19 | 0.99 ± 0.52 | 0.255 |
| US - EC Right (deg/sec) | 2.31 ± 0.59 | 2.23 ± 0.77 | 0.715 |
| LOS - RT Forward (sec) | 0.89 ± 0.34 | 0.92 ± 0.22 | 0.763 |
| LOS - RT Right (sec) | 0.83 ± 0.32 | 0.90 ± 0.24 | 0.422 |
| LOS - RT Back (sec) | 0.81 ± 0.25 | 0.78 ± 0.32 | 0.794 |
| LOS - RT Left (sec) | 0.86 ± 0.25 | 0.84 ± 0.28 | 0.83 |
| LOS - MVL Forward (deg/sec) | 3.64 ± 1.48 | 3.76 ± 1.54 | 0.738 |
| LOS - MVL Right (deg/sec) | 5.29 ± 2.88 | 4.56 ± 2.33 | 0.222 |
| LOS - MVL Back (deg/sec) | 3.01 ± 1.14 | 3.06 ± 1.09 | 0.825 |
| LOS - MVL Left (deg/sec) | 4.92 ± 2.29 | 5 ± 1.79 | 0.867 |

mCTIB, Modified Clinical Test of Sensory Interaction on Balance; US, Unilateral Stance; EO, Eyes open; EC, Eyes closed; LOS, Limits of Stability; RT, Reaction time; MVL, Movement velocity.

Table 3. Comparison of balance scores in non-fatigue group.

| Variables | Pretest Mean ± SD | Posttest Mean ± SD | T test |
|-----------------------------|----------------------|-----------------------|--------|
| | | | P |
| mCTIB Firm - EO (deg/sec) | 0.37 ± 0.13 | 0.38 ± 0.23 | 0.885 |
| mCTIB Firm - EC (deg/sec) | 0.35 ± 0.14 | 0.35 ± 0.21 | 1 |
| mCTIB Foam - EO (deg/sec) | 0.66 ± 0.11 | 0.67 ± 0.16 | 1 |
| mCTIB Foam - EC (deg/sec) | 1.40 ± 0.48 | 1.31 ± 0.31 | 0.329 |
| US - EO Left (deg/sec) | 1.24 ± 0.76 | 1.01 ± 0.50 | 0.093 |
| US - EC Left (deg/sec) | 2.87 ± 1.03 | 2.79 ± 1.12 | 0.761 |
| US - EO Right (deg/sec) | 1.13 ± 0.70 | 1.36 ± 1.04 | 0.222 |
| US - EC Right (deg/sec) | 2.81 ± 1.13 | 2.72 ± 1.33 | 0.647 |
| LOS - RT Forward (sec) | 0.91 ± 0.38 | 0.88 ± 0.20 | 0.742 |
| LOS - RT Right (sec) | 0.80 ± 0.29 | 0.86 ± 0.30 | 0.343 |
| LOS - RT Back (sec) | 0.82 ± 0.29 | 0.72 ± 0.26 | 0.139 |
| LOS - RT Left (sec) | 0.86 ± 0.23 | 0.71 ± 0.29 | 0.068 |
| LOS - MVL Forward (deg/sec) | 3.29 ± 1.23 | 3.83 ± 1.36 | 0.14 |
| LOS - MVL Right (deg/sec) | 4.16 ± 1.55 | 5.02 ± 2.38 | 0.061 |
| LOS - MVL Back (deg/sec) | 3 ± 0.97 | 3.31 ± 0.97 | 0.182 |
| LOS - MVL Left (deg/sec) | 5.03 ± 1.84 | 5.99 ± 2.49 | 0.118 |

mCTIB, Modified Clinical Test of Sensory Interaction on Balance; US, Unilateral Stance; EO, Eyes open; EC, Eyes closed; LOS, Limits of Stability; RT, Reaction time; MVL, Movement velocity.

fatigue on balance scores (mCTSIB, US, and LOS) showed no significant difference in all tested positions ($p>0.05$) (Table 2). The comparison of balance scores (mCTSIB, US, and LOS) from pretest to posttest in the Non-Fatigue Group showed no significant difference in all the tested positions

(Table 3). The posttest values were used to compare the differences of balance scores between the three groups. The comparison of balance scores (mCTSIB, US, and LOS) between Quadriceps, Calf and Non-Fatigue groups showed no significant difference (Table 4).

Table 4. Comparison of balance scores among quadriceps, calf and non-fatigue groups.

| Variable | Quadriceps Group Mean ± SD | Calf Group Mean ± SD | Non Fatigue Group Mean ± SD | ANOVA P |
|-----------------------------|-------------------------------|-------------------------|--------------------------------|------------|
| mCTIB Firm - EO (deg/sec) | 0.45 ± 0.15 | 0.44 ± 0.13 | 0.380 ± 0.24 | 0.493 |
| mCTIB Firm - EC (deg/sec) | 0.35 ± 0.09 | 0.50 ± 0.50 | 0.353 ± 0.22 | 0.345 |
| mCTIB Foam - EO (deg/sec) | 0.77 ± 0.21 | 0.68 ± 0.11 | 0.67 ± 0.16 | 0.164 |
| mCTIB Foam - EC (deg/sec) | 1.33 ± 0.39 | 1.25 ± 0.36 | 1.31 ± 0.31 | 0.846 |
| US - EO Left (deg/sec) | 1.16 ± 0.58 | 0.81 ± 0.09 | 1.01 ± 0.50 | 0.107 |
| US - EC Left (deg/sec) | 2.80 ± 0.83 | 2.13 ± 0.91 | 2.79 ± 1.12 | 0.100 |
| US - EO Right (deg/sec) | 1.17 ± 0.39 | 0.99 ± 0.53 | 1.36 ± 1.04 | 0.378 |
| US - EC Right (deg/sec) | 3.06 ± 0.93 | 2.23 ± 0.77 | 2.72 ± 1.33 | 0.103 |
| LOS - RT Forward (sec) | 0.88 ± 0.34 | 0.92 ± 0.22 | 0.88 ± 0.20 | 0.900 |
| LOS - RT Right (sec) | 0.85 ± 0.32 | 0.90 ± 0.25 | 0.85 ± 0.30 | 0.853 |
| LOS - RT Back (sec) | 0.78 ± 0.33 | 0.78 ± 0.32 | 0.71 ± 0.26 | 0.827 |
| LOS - RT Left (sec) | 0.92 ± 0.24 | 0.84 ± 0.28 | 0.71 ± 0.29 | 0.113 |
| LOS - MVL Forward (deg/sec) | 3.44 ± 1.62 | 3.76 ± 1.55 | 3.83 ± 1.37 | 0.758 |
| LOS - MVL Right (deg/sec) | 4.60 ± 1.48 | 4.57 ± 2.33 | 5.02 ± 2.38 | 0.807 |
| LOS - MVL Back (deg/sec) | 2.89 ± 0.84 | 3.06 ± 1.10 | 3.31 ± 0.98 | 0.500 |
| LOS - MVL Left (deg/sec) | 5.06 ± 1.73 | 5.00 ± 1.80 | 5.99 ± 2.5 | 0.335 |

mCTIB, Modified Clinical Test of Sensory Interaction on Balance; US, Unilateral Stance; EO, Eyes open; EC, Eyes closed; LOS, Limits of Stability; RT, Reaction time; MVL, Movement velocity.

Discussion

The aims of this study were to investigate and compare the effects of quadriceps and calf muscles fatigue on quiet standing balance in young healthy adult males. In the current study, participants were tested for sway velocity (degrees per second) for mCTSIB and US under four conditions. In addition, limits of stability in eight directions were measured for the movement velocity (degrees per second) and reaction time (sec). The results showed no significant difference between pretest and posttest scores for all the variables. Our findings are not in agreement with the previous studies, which reported significant effects of quadriceps and calf muscles fatigue on quiet standing balance^{25,35}. Another study reported reduced dynamic balance following muscle fatigue in adult male³⁶. Similarly, other studies reported reduced dynamic balance following muscle fatigue^{37,38}. This difference in the results might be due to various methodological differences between the present and previous studies. The previous studies have investigated older participants whereas in our study, we used young participants^{25,35}. Many studies have investigated the effects of aging on balance. Zettel et al.³⁹ concluded that the attentional demand related with balance recovery was impaired in older adults. Abdulvahabi et al.³⁵ and Islami et al.²⁵ reported that there was a meaningful decrease in balance score posttest (after fatigue) as compared with pretest (baseline) conditions in elderly subjects, which indicated that aging had a negative effect on dynamic standing balance. In contrast, our participants were comprised of young adults and we found that fatigue did not have a significant effect

on standing balance. In addition, the gender of participants in this study was males while previous studies that reported a significant difference in the effect of fatigue on balance studied female participants^{25,35}. This might be due to differences in body muscle mass between genders. Studies have shown that males have greater skeletal muscle mass than females. In one study that examined 468 males and females, they determined that males had an average of 33 kilograms of muscle compared to the 21 kilograms found in females. The males had 40 percent more muscle mass in the upper body and 33 percent more in the lower body⁴⁰. Higher muscle mass might contribute to better standing balance for males compared to females.

Another factor can be related to the compensatory mechanisms thought to come into play after implementation of a fatigue protocol for maintaining standing balance. The fatigue of localized muscle does not affect standing balance because of compensatory mechanisms. Other muscle groups might work together in order to compensate for the fatigued muscle group. Therefore, no effect of fatigue would appear on general standing balance^{25,35}. There were methodological differences used in detecting standing balance changes amongst the various studies^{25,35}. In the present study, we used the Balance Master, because of its ability to detect minimal changes during static and dynamic standing balance^{25,35}. Previous reports used functional balance measures such as the Berg Balance Scale, which combines static and dynamic measures into one composite score and might not necessarily reflect the individual static and dynamic standing performances²⁵. Another study used the Star Excursion test,

which focuses on dynamic balance only, measured manually in centimeters, and is not as accurate as the computerized measures from the Balance Master³⁵.

In the present study, we reported no significant difference in posttests balance scores for all the groups. In contrast, previous studies reported significant differences between quadriceps and calf muscle fatigue on balance^{25,35}. In addition, both quadriceps and calf muscles contain two types of muscle fibers: slow twitch (type I) and fast twitch (type II). A previous study reported that the soleus, gastrocnemius, and vastus lateralis muscles had a high proportion of slow twitch fibers (Type I)⁴¹. In addition, the glycolytic activity was lower in the slow twitch fibers (Type I) compared to those with a high proportion of high twitch fibers (Type II)⁴¹. Since, the antigravity muscles in the lower limb such as quadriceps and calf muscles have more type I fibers; therefore, these muscles may have a higher resistance to fatigue than other muscles. Since both the muscles are similar in morphological structure, it is assumed that the fatigue of both muscles (quadriceps and calf muscles) have similar effects on balance^{41,42}.

This study is limited to healthy young adult subjects with no history of falls or balance impairments. To what degree acute fatigue would affect balance in young subjects with a history of falling is unknown and worthy of study. In addition, the present study assessed the participant's perceived level of fatigue but the lasting effects of fatigue were not tested. These could also affect the present results.

Future studies should incorporate EMG data of lower limb musculature in order to a better understanding the possible compensatory measures exhibited by our subjects under fatigued conditions. In addition, using surface EMG is highly recommended to ensure that the participants reached a fatigued state following the fatiguing protocol, and a threshold for decreased in muscular strength could be established. In addition, the effects of physiological parameters of fatigued muscles such as oxidative free radicals, lactate dehydrogenase, and creatine kinase on balance performance could be studied.

In conclusion, the fatigue of the quadriceps or calf muscles did not influence standing balance in healthy young adult males. Future longitudinal studies are warranted to further understanding the mechanisms behind localized muscle fatigue effects on standing balance in subjects of different age groups of both genders.

Acknowledgement

The authors would like to extend their sincere appreciation to the Deanship of Scientific Research at King Saud University for funding this research through the research group NO. RGP-VPP-209.

Ethics approval and consent to participate

The study was approved by the ethics committee of the College of Applied Medical Sciences, King Saud University. The participants were asked to sign a written informed consent form approved by the institution ethics committee of King Saud University.

Funding

This project was funded by the Deanship of Scientific Research, King Saud University through the research group NO. RGP-VPP-209. The funding body played no role in the study design, manuscript writing, or decision to submit the manuscript for publication.

Authors' contributions

AAJ: participated in the study design, participated in the data collection, and drafted the manuscript. SA: corresponding author, participated in the study design, participated in the data collection, revised the manuscript critically, and finalized the manuscript. AA: participated in the study design, helped with the ethics application and revised the manuscript critically. HZ: participated in the study design, developed the protocol, and revised the manuscript critically. EA: participated in the study design and revised the manuscript critically. SME: participated in the study design and revised the manuscript critically. CT: participated in the study design and revised the manuscript critically. All authors read and approved the final manuscript.

References

1. Aslan UB, Cavlak U, Yagci N, y Akdag B. Balance performance, aging and falling: A comparative study based on a Turkish sample. *Arch Gerontol Geriatr* 2008;46:283-292.
2. Odebiyi DO, Oderinde MO, Olaogun MO. Balance, gait and fear of falling in the elderly-a pilot study in community-based rehabilitation homes. *West Afr J Med* 2008;27:245-249.
3. Russell KA, Palmieri RM, Zinder SM, Ingersoll CD. Sex differences in valgus knee angle during a single-leg drop jump. *J Athl Train* 2006;41:166-171.
4. Gribble PA, Hertel J. Effect of hip and ankle muscle fatigue on unipedal postural control. *J Electromyogr Kinesiol* 2004;14:641-646.
5. Rubenstein LZ. Falls in older people: epidemiology, risk factors and strategies for prevention. *Age Ageing* 2006;35:37-41.
6. Bellew JW, Fenter PC. Control of balance differs after knee or ankle fatigue in older women. *Arch Phys Med Rehabil* 2006;87:1486-1489.
7. Nam HS, Park DS, Kim DH, Kang HJ, Lee DH, Lee SH, Her JG, Woo JH, Choi SY. The relationship between muscle fatigue and balance in the elderly. *Ann Rehabil Med* 2013;37:389-395.
8. Paillard T. Effects of general and local fatigue on postural control: a review. *Neurosci Biobehav Rev* 2012;36:162-176.
9. James CR, Scheuermann BW, Smith MP. Effects of two neuromuscular fatigue protocols on landing performance. *J Electromyogr Kinesiol* 2010;20:667-675.
10. Enoka RM, Duchateau J. Muscle fatigue: what, why and how it influences muscle function. *J Physiol* 2008; 586:11-23.
11. Al-Mulla MR, Sepulveda F, Colley M. A review of non-invasive techniques to detect and predict localised muscle fatigue. *Sensors (Basel)* 2011;11:3545-3594.

12. Bisson EJ, Remaud A, Boyas S, Lajoie Y, Bilodeau M. Effects of fatiguing isometric and isokinetic ankle exercises on postural control while standing on firm and compliant surfaces. *J Neuroeng Rehabil* 2012;9:39.
13. Zemková E, Hamar D. The agility test in functional diagnostics of athletes. *Acta Univ Palacki Olomuc, Gymn* 2004;34:2.
14. Trudelle-Jackson EJ, Jackson AW, Morrow J. Muscle strength and postural stability in healthy, older women: implications for fall prevention. *J Phys Act Health* 2006; 3:292.
15. Kavanagh JJ, Morrison S, Barrett RS. Lumbar and cervical erector spinae fatigue elicit compensatory postural responses to assist in maintaining head stability during walking. *J Appl Physiol* 2006;101:1118-1126.
16. Kent-Braun JA. Skeletal Muscle Fatigue in Old Age: Whose Advantage? *Exerc Sport Sci Rev* 2009;37:3-9.
17. Cetin N, Bayramoglu M, Aytar A, Surenkok O, Yemisci OU. Effects of Lower Extremity Muscle Fatigue on Balance. *The Open Sports of Medicine Journal* 2008;2:11-17.
18. Lanza IR, Larsen RG, Kent-Braun JA. Effects of old age on human skeletal muscle energetics during fatiguing contractions with and without blood flow. *J Physiol* 2007;583:1093-1105.
19. Mademli L, Arampatzis A. Mechanical and morphological properties of the triceps surae muscle-tendon unit in old and young adults and their interaction with a submaximal fatiguing contraction. *J Electromyogr Kinesiol* 2008;18:89-98.
20. Abd-Elfattah HM, Abdelazeim FH, Elshennawy S. Physical and cognitive consequences of fatigue: A review. *J Adv Res* 2015;6:351-358.
21. Hunter SK, Duchateau J, Enoka RM. Muscle fatigue and the mechanisms of task failure. *Exerc Sport Sci Rev* 2004;32:44-49.
22. Dideriksen JL, Enoka RM, Farina D. Neuromuscular adjustments that constrain submaximal EMG amplitude at task failure of sustained isometric contractions. *J Appl Physiol* 2011;111:485-494.
23. Gear WS. Effect of different levels of localized muscle fatigue on knee position sense. *J Sports Sci Med* 2011; 10:725-730. eCollection 2011.
24. Gribble PA, Hertel J. Effect of lower-extremity muscle fatigue on postural control. *Arch Phys Med Rehabil* 2004;85:589-592.
25. Islami F, Fallah Z, Mahdavi S, Hesari AF, Janani H. Effect of quadriceps and ankle plantar flexor muscle fatigue on balance of elderly women. *HealthMed* 2012;6:875-878.
26. Vuillerme N, Forestier N, Nougier V. Attentional demands and postural sway: the effect of the calf muscles fatigue. *Med Sci Sports Exerc* 2002;34:1907-1912.
27. Susco TM, Valovich Mcleod TC, Gansneder BM, Shultz SJ. Balance Recovers Within 20 Minutes After Exertion as Measured by the Balance Error Scoring System. *J Athl Train* 2004;39:241-246.
28. Vuillerme N, Pinsault N, Vaillant J. Postural control during quiet standing following cervical muscular fatigue: effects of changes in sensory inputs. *Neurosci Lett* 2005;378:135-139.
29. Ledin T, Fransson PA, Magnusson M. Effects of postural disturbances with fatigued triceps surae muscles or with 20% additional body weight. *Gait Posture* 2004;19:184-193.
30. Vuillerme N, Burdet C, Isableu B, Demetz S. The magnitude of the effect of calf muscles fatigue on postural control during bipedal quiet standing with vision depends on the eye-visual target distance. *Gait Posture* 2006;24:169-172.
31. Cohen H, Blatchly CA, Gombash LL. A study of the clinical test of sensory interaction and balance. *Phys Ther* 1993;73:346-51.
32. Wrisley DM, Whitney SL. The effect of foot position on the modified clinical test of sensory interaction and balance1. *Arch Phys Med Rehabil* 2004;85:335-338.
33. Ageberg E, Roberts D, Holmström E, Friden T. Balance in single-limb stance in healthy subjects-reliability of testing procedure and the effect of short-duration submaximal cycling. *BMC Musculoskelet Disord* 2003;4:14.
34. Zouita Ben Moussa A, Zouita S, Dziri C, Ben Salah FZ. Single-leg assessment of postural stability and knee functional outcome two years after anterior cruciate ligament reconstruction. *Ann Phys Rehabil Med* 2009; 52:475-484.
35. Abdolvahabi Z, Bonab SS, Rahmati H, Naini SS. The Effects of Ankle Plantar Flexor and Knee Extensor Muscles Fatigue on Dynamic Balance of the Female Elderly. *World Applied Sciences Journal* 2011; 15:1239-1245.
36. Zulfikri N, Justine M. Effects of Kinesio® Taping on Dynamic Balance Following Fatigue: a Randomized Controlled Trial. *Phys Ther Res* 2017;20:16-22.
37. Gribble PA, Hertel J, Plisky P. Using the Star Excursion Balance Test to assess dynamic postural-control deficits and outcomes in lower extremity injury: a literature and systematic review. *J Athl Train* 2012;47:339-357.
38. Steib S, Zech A, Hentschke C, Pfeifer K. Fatigue-induced alterations of static and dynamic postural control in athletes with a history of ankle sprain. *J Athl Train* 2013; 48:203-208.
39. Zettel JL, McIlroy WE, Maki BE. Effect of competing attentional demands on perturbation-evoked stepping reactions and associated gaze behavior in young and older adults. *J Gerontol A Biol Sci Med Sci* 2008; 63:1370-1379.
40. Janssen I, Heymsfield SB, Wang ZM, Ross R. Skeletal muscle mass and distribution in 468 men and women aged 18-88 yr. *J Appl Physiol* 2000;89:81-88.
41. Gollnick PD, Sjödin B, Karlsson J, Jansson E, Saltin B. Human soleus muscle: a comparison of fiber composition and enzyme activities with other leg muscles. *Pflugers Arch* 1974;348:247-255.
42. Esbjörnsson-Liljedahl M, Sundberg CJ, Norman B, Jansson E. Metabolic response in type I and type II muscle fibers during a 30-s cycle sprint in men and women. *J Appl Physiol* 1999;87:1326-1332.