

Alendronate as an effective countermeasure to disuse induced bone loss

A.D. LeBlanc¹, T.B. Driscoll¹, L.C. Shackelford², H.J. Evans³,
N.J. Rianon¹, S.M. Smith², D.L. Feedback², D. Lai⁴

¹Department of Medicine, Baylor College of Medicine, ²Musculoskeletal Laboratory, Johnson Space Center, NASA, ³Life Sciences, Systems and Services, Wyle Laboratories, ⁴School of Public Health, University of Texas, Houston, Texas, USA

Abstract

Microgravity, similar to disuse immobilization on earth, causes rapid bone loss. This loss is believed to be an adaptive response to the reduced musculoskeletal forces in space and occurs gradually enough that changes occurring during short duration space flight are not a concern. Bone loss, however, will be a major impediment for long duration missions if effective countermeasures are not developed and implemented. Bed rest is used to simulate the reduced mechanical forces in humans and was used to test the hypothesis that oral alendronate would reduce the effects of long duration (17 weeks) inactivity on bone. Eight male subjects were given daily oral doses of alendronate during 17 weeks of horizontal bed rest and compared with 13 male control subjects not given the drug. Efficacy was evaluated based on measurements of bone markers, calcium balance and bone density performed before, during and after the bed rest. The results show that oral alendronate attenuates most of the characteristic changes in bone that are associated with long duration bed rest and presumably space flight.

Keywords: Microgravity, Bone, Bed Rest, Alendronate

Introduction

The body's normal adaptive response to reduced gravitational forces causes untoward effects in many organ systems, e.g., neurovestibular disturbances, cardiovascular deconditioning, loss of red blood cells and plasma volume, loss of muscle mass and strength and decreased bone density¹. The potential for significant and possibly irreversible loss of bone is one of the most important medical concerns for long-duration manned space flight. Despite the implementation of a comprehensive exercise program aboard the Mir space station, regional decreases in bone density of 1-1.6% per month occur²⁻⁴. This bone loss is not uniform, displaying significant heterogeneity between individuals and between bone sites within a given individual. For example, generally there are no losses in the arms while more than 97% of the total body bone mineral loss originates from the pelvis and lower limbs⁴. This regional variation probably reflects the

change in gravitational and muscular forces experienced by the skeleton during routine ambulation and exercising in one G compared to microgravity. While the data is currently being analyzed and not yet published, it appears that recovery does occur, although slowly and sometimes incompletely. Similar to the variable response seen during the loss of bone, recovery of bone appears to show variability in the rate and degree of bone restitution following flight⁵.

NASA's ability to assure the health of the astronauts during long-duration space flight is compromised without an effective countermeasure to disuse bone loss. Countermeasures for disuse bone loss could employ exercise, biochemical prescriptions or some combination of both. Exercise is the most desirable countermeasure for maintaining muscle volume and strength since it has positive effects on the cardiovascular as well as the musculoskeletal system. The results from Mir, however, indicate that exercise alone may prove difficult as the only countermeasure for bone loss prevention, i.e., crew members continued to lose bone. As opposed to exercise, an effective biochemical intervention would require minimal crew time, minimal storage space and no power. No biochemical countermeasure has been attempted during weightlessness. However, several attempts

Corresponding author: Adrian LeBlanc, PhD, Baylor College of Medicine, 6550 Fannin St., Suite 1260, Houston, TX 77030, USA
E-mail: aleblanc@bcm.tmc.edu

Accepted 5 March 2002

have been made using ground-based bed rest studies. Bed rest is commonly used to simulate some of the effects of space flight on the musculoskeletal system. The skeletal changes, i.e., decreases in bone density, increased bone resorption markers, increased urinary and fecal calcium (Ca) excretion and negative Ca balance were found to be qualitatively similar, although smaller in magnitude, during bed rest than in space flight²⁻⁸.

Two of the early bisphosphonates, etidronate and clodronate, were tested as potential countermeasures to disuse bone loss. During the last eight weeks of a 20-week bed rest study, Ca balance shifted toward positive with high dose etidronate (20 mg/kg/day). Significant calcaneal mineral losses however were observed⁹⁻¹⁰. This drug was not pursued since etidronate is associated with an accumulation of osteoid tissue in both animals and man when it was given at the anti-resorptive dose for extended periods of time¹¹⁻¹². Clodronate was tested in an unpublished 17-week bed rest study, the results of which indicated that Ca balance was maintained. CT-densitometry of the spine in the nine male treated subjects approached statistical significance in preventing lumbar spine bone loss compared to five controls. Single photon absorptiometry of the calcaneus, however, showed no statistically significant difference between the two groups and one treated test subject showed severe calcaneal density loss while maintaining normal Ca balance¹³. Clodronate was withdrawn from clinical investigation in the United States due to a potential serious adverse reaction. In 1995, the FDA approved alendronate for the treatment of postmenopausal osteoporosis and Paget's disease of bone. One published short duration bed rest study indicated that alendronate might be safe and effective in preventing bed rest induced bone loss¹⁴. This study compared two groups of men, one treated with alendronate (n=8) and the other with placebo (n=8) and consisted of 2 weeks of pre-bed rest and 3 weeks of bed rest. The treated subjects were given 20 mg/day for all 5 weeks. During the bed rest phase subjects sat up briefly to ingest the pill and were then returned to the lying down position. During the pre-bed rest ambulatory control period, alendronate-treated subjects exhibited reduced urinary calcium compared to the placebo group. Both the placebo and alendronate groups showed significant increases in urinary calcium excretion during bed rest. The excretion during bed rest in the alendronate group, however, was below the baseline (pre-bed rest) level of the placebo group, suggesting that this dose level might be effective against bone loss. During the ambulatory phase of the study, two of the treated subjects had asymptomatic decreased serum Ca that reversed during bed rest. No adverse reactions to alendronate were noted in this study.

Although the above studies suggest that bisphosphonates might be effective, it was important to NASA physicians to verify safety and efficacy using ground-based models prior to

routine use on long duration space missions. The purpose of this study was to test the hypothesis that oral alendronate (10mg/day) would reduce the effects of long duration (17 weeks) bed rest on bone as evidenced by the effects on bone markers, calcium balance and bone density. Because the drug was not approved for use in young women, this study was limited by NASA to normal males in an age group similar to the astronaut corps. The alendronate treated subjects (n=8) were compared to untreated bed rest male subjects (n=13). Since the drug was not expected to have an effect on muscle, changes in lean body mass (LBM) were used to determine comparability of the treated and control groups.

Methods

The Baylor College of Medicine and the NASA, Johnson Space Center Human Use Committees reviewed and approved this research protocol. Informed consent was obtained from all subjects. The design of this study was to evaluate drug efficacy using each individual as his own control (pre vs. bed rest) and by comparison with untreated male controls. The latter consisted of 8 men from previous 17-week studies by the same investigative team using the same research protocol⁷⁻⁸ and five men from a concurrent bed rest study investigating another countermeasure.

Research subjects were recruited using local newspaper advertisements. All potential subjects were required to: pass a military Class III physical (ENT exam, heart sounds, blood pressure, EKG, blood count, blood and urine analysis and drug screen), have a bone mineral density (BMD) result within two standard deviations above or below the age/sex-matched mean and normal drug and psychological (MMPI) screens. Individuals with a history of upper G.I. problems such as ulcers, dysphagia, symptomatic esophageal disease, gastritis, or duodenitis were excluded from the study. In addition, the treated subjects were tested for pill tolerance. The tolerance test consisted of ingesting a 10 mg dose of alendronate (FOSAMAX, Merck & Co) with 8 ounces of water and then immediately lying down for 30 minutes. No tested subject reported any discomfort from this procedure.

A total of nine men were recruited to participate in the drug treatment portion of the study. One individual passed all the pre-study physicals and screening procedures, but was subsequently found to have a hiatal hernia. This individual was removed from the study resulting in a total of eight males in the treatment group. One other subject, early in the bed rest period, complained of heartburn in the evening and revealed having similar problems prior to entering the study. The symptoms and their timing, occurring in the evening, did not seem related to the ingestion of alendronate, taken in the morning. Therefore, the subject was put on 50 mg/day Prilosec and continued in the study without further problems. The eight treated subjects were 22-56 years of age

(mean=35), 160-185 cm in height (mean=173) and 65-98 kg in weight (mean=77). The controls consisted of 13 men, 22-56 years of age (mean=32), 156-191 cm in height (mean=171) and 65-99 kg in weight (mean=76). The protocol for all subjects was the same except for some differences in measured variables between previous and current controls as indicated in the results.

The studies were divided into three periods; a pre-bed rest period, 17 weeks of bed rest and a post-bed rest recovery period. During the pre-bed rest period, the subjects were ambulatory while housed in the testing facility. During this time, all baseline measurements were obtained. In bed rest, the subjects were not permitted to sit up, but were allowed to raise themselves on one elbow while maintaining continuous contact with the bed, for eating, using a portable computer or reading. For defecation, the subjects were allowed to move to a portable commode, located next to the bed. Generally this occurred about 3-7 times per week and required prior permission from ward personnel. The subjects were closely monitored at all times to prevent unauthorized activity. The treated subjects were given a 10 mg tablet of alendronate each day during the bed rest period. The fasting subjects were asked to briefly sit up each morning to take the pill with 8 ounces of water. After the pill was swallowed, subjects immediately returned to lying down. Subjects consumed no food, medicine or drink (other than plain water) for 12 hours before and 90 minutes after ingestion of the alendronate dose. During the post-bed rest period, the subjects were allowed to resume ambulatory activities *ad lib*, but were cautioned to avoid excessive walking during the first several days.

Subjects were maintained on a metabolic diet during the entire study. The diet was prepared in a similar fashion as previously described⁷. Briefly, the diet was composed of 7 daily menus, each consisting of three meals and an evening snack. All food portions were separately weighed. The subjects were required to eat all food. On average, the diet contained about 1 g of calcium, 1.7 g of phosphorus, and 15 g of nitrogen per day. In addition, all subjects received a hexavitamin tablet once daily which provided approximately 14,000 I.U. of vitamin A, 630 I.U. of vitamin D, 3 mg thiamin, 5 mg riboflavin, 42 mg niacin, and 192 mg ascorbic acid. The mineral content of the diet was confirmed on 20 different weeks during the study. During these weeks additional meals were prepared that were subsequently homogenized, ashed and analyzed. A description of the mineral analysis has been published previously¹⁵. The subjects were weighed weekly using an in-bed scale. The caloric intake was adjusted to maintain approximate constant body weight during the study. In these instances, the caloric content was adjusted while maintaining a constant mineral content using a diet analysis software package (Nutritionist V, First Databank).

Bone densitometry

Bone mineral of the whole body, spine (L1-L4), femur neck, trochanter, proximal (1/3 site) and distal radius and calcaneus were measured 3 times during baseline ambulation and 3 times at the end of bed rest. The mean pre- and post-bed rest values were used for analysis. All current bone densitometry measurements were obtained using a Hologic QDR 2000+ whole body densitometer. The older published BMD measurements were obtained from various DEXA instruments, but all pre- and post- measurements on a given individual were performed on the same instrument and obtained and analyzed by the same operator. Routine acquisition and analysis software were used to obtain BMD and soft tissue scans except for the calcaneus. These scans were obtained using a special jig and the Hologic spine acquisition software. Subregions of the whole body were those defined by Hologic software and included the pelvis, legs (all of both limbs and feet), arms (both limbs and hands). Phantom scans were performed regularly in order to assess the short-term and long-term performance of the densitometer. Approximately 20 times per month (and on the morning of each day of testing), a Hologic spine phantom was scanned. This phantom is an anatomical model of the lumbar spine, composed of hydroxyapatite and encased in plastic. The precision error for phantom bone mineral density is approximately 0.5%. The precision of bone and soft tissue DEXA measurements in our laboratory has been published¹⁶⁻¹⁸. The most recent laboratory BMD precision measurements expressed as %CV are: lumbar spine, 1.4%; femur neck, 1.5%; trochanter, 1.8%; ultra distal radius, 2.1%; whole body, 0.6%.

Urine and fecal Ca excretion and Ca balance

Twenty-four hour urine samples were collected daily throughout the study. During collection, the 24 hour voidings were stored at 4°C. At the end of each 24 hour period, the total volume was measured and duplicate aliquots (2% of total volume) were obtained. One aliquot was acidified with HCl and both aliquots were then frozen. The aliquots from each day were combined to form a 7 day pool and stored frozen at -22°C in preparation for later Ca analysis. Individually collected fecal samples were combined into weekly, 7 day pools and stored frozen for subsequent Ca analysis. Urinary Ca was analyzed weekly using plasma-mass spectrometry, while fecal Ca was analyzed on 4 occasions; during 3 weeks of baseline, and at 4-6 weeks, 10-12 weeks and 15-17 weeks of bed rest. Ca balance was determined using fecal and urinary Ca data during these same intervals. Calculations of metabolic balance were based on the following: Ca intake in mg/day minus output mg/day (urinary + fecal Ca). Net balance was obtained as the difference between baseline and bed rest.

Markers of bone metabolism and regulatory hormones

In order to determine changes in the markers of bone metabolism and regulatory hormones, blood and urine samples were obtained during the ambulatory period and the 3 bed rest time periods described above. The blood samples were obtained in the early morning, after a 12 hour fast, during the 3rd day of the last week of each 3-week interval. Measures obtained from serum were: total calcium (ion-sensitive electrode, Beckman Coulter Instruments, Brea, CA); ionized Ca (ion-sensitive electrode, i-STAT Corporation, Princeton, NJ); total alkaline phosphatase (spectrophotometry); bone-specific alkaline phosphatase (enzyme-linked immunosorbent assay, Metra Biosystems, Palo Alto, CA); 1,25 Dihydroxyvitamin D and 25 Hydroxyvitamin D by radioimmunoassay (DiaSorin, Inc., Stillwater, MN); osteocalcin (radioimmunoassay, Biomedical Technologies, Stoughton, MA); and mid-molecule and intact molecule PTH (radioimmunoassay, Nichols Institute Diagnostics, San Juan Capistrano, CA). Measures obtained from urine were: total calcium (ion-sensitive electrode, Beckman Coulter Instruments, Brea, CA); n-telopeptide (enzyme-linked immunosorbent assay, Ostex Int., Seattle, WA); and pyridinium and deoxypyridinoline crosslinks (enzyme-linked immunosorbent assay, Metra Biosystems, Palo Alto, CA). The bone specific alkaline phosphatase, intact molecule PTH and 25(OH)D were obtained on the concurrent controls and drug treated subjects only.

Lean body mass measurements

Changes in whole body and regional lean body mass were determined from the whole body DEXA scans. Reproducibility of these measurements in our laboratory has been determined in a published precision study¹⁸. Precision values for LBM are: arms, 1.8%; legs, 1.3%; whole body, 0.9%.

Statistics

Except for the Ca excretion given in mg/day, the data are expressed as percentage change from pre-bed rest values. The percentage changes are reported as mean \pm SE for the control and treatment groups. A paired t-test was used to determine if the changes during bed rest for the individual variables were statistically significant. Differences between the control and treatment group means were tested using unpaired t-test. Within and between group differences for groups of variables were tested using Wilk's Lambda statistic¹⁹ by SAS proc GLM (SAS Institute, NC). The groups tested using the latter were: BMD, urine and fecal Ca excretion, resorption markers, formation markers and lean body mass. A p value of less than 0.05 was used for determining significance.

Results

Table 1 gives the BMD changes for the control and treated subjects. During bed rest, the BMD of the control group decreased in all regions except the arms and radius. The arms and radius were unchanged as reported previously⁷. Only the 5 concurrent control data are presented for the radius since the instrument (Osteon, Inc.) used previously, scanned somewhat different regions and therefore could not be combined with the current DEXA data. The results, however, were the same. The treated group showed no loss in BMD except the calcaneus which decreased about 5% compared to 10% in the controls. As opposed to a loss in the control group, the femur neck of the treated group showed an increase in BMD. Comparing the treated subjects to controls, significant differences were found in the spine, femur neck, trochanter, and pelvis. Compared to controls, the changes in the whole body, legs and arms although smaller in the treated group, did not reach significance. All BMD regions, from Table 1, that decreased in the control group (lumbar spine, femoral neck, trochanter, calcaneus, total body, pelvis, legs) were combined for testing for a treatment effect using the Wilk's Lambda statistic. For the controls, the within group differences were significant ($p=0.006$) while the treatment group were not ($p=0.68$), i.e., the controls lost BMD while the treated subjects did not. The between group testing was significant ($p=0.019$), demonstrating a significant effect from the treatment.

Table 2 gives the hormonal and bone marker changes as a percentage of baseline for the control and treated subjects. In the controls, the three resorption markers all increased and the PTH (intact molecule) decreased during bed rest. In the current study both intact and mid-molecule PTH assays were performed while previously, only the latter was performed. In the treated group, the NTx resorption marker decreased while the other two resorption markers increased, but significantly less than in the controls. The formation markers all decreased relative to baseline in the treated group while unchanged in the controls. Figures 1-3 show the mean percentage change in NTx, osteocalcin and intact molecule PTH for the control and treated groups for the 3 measured time points during bed rest. Within group changes in the control and treatment groups of the combined resorption markers using the Wilk's Lambda statistic were significant ($p<0.001$). Comparing the resorption markers between the two groups, the Wilk's Lambda was significant ($p<0.001$), confirming a significant drug effect. Similar to the paired t-test results, the Wilk's Lambda indicated no significant within group change in the formation markers in the controls, but a significant decrease in the treated group ($p<0.001$). Unlike the unpaired t-test, the Wilk's Lambda did not show a significant difference in formation markers between the control and treated groups. Both serum Ca and

| | Control (n=13) | | | Alendronate (n=8) | | |
|-----------------|----------------|---------------|--------------------------|-------------------|---------------|--------------------------|
| | Baseline | Post-Bed Rest | % Change | Baseline | Post-Bed Rest | % Change |
| Lumbar spine | 1.11 ± 0.04 | 1.09 ± 0.03 | -1.6 ± 0.8 ^a | 1.09 ± 0.06 | 1.11 ± 0.06 | 1.8 ± 0.9 ^c |
| Femoral neck | 0.93 ± 0.04 | 0.91 ± 0.04 | -2.0 ± 0.9 ^a | 1.00 ± 0.09 | 1.02 ± 0.09 | 2.1 ± 0.6 ^{b,d} |
| Trochanter | 0.77 ± 0.02 | 0.74 ± 0.03 | -3.9 ± 0.6 ^b | 0.80 ± 0.06 | 0.79 ± 0.06 | 0.0 ± 0.7 ^d |
| Calcaneus | 0.58 ± 0.04 | 0.52 ± 0.04 | -10.3 ± 2.3 ^b | 0.72 ± 0.04 | 0.68 ± 0.03 | -4.9 ± 1.6 ^a |
| Distal radius | 0.55 ± 0.05* | 0.55 ± 0.04* | -0.5 ± 0.6* | 0.56 ± 0.02 | 0.57 ± 0.02 | 0.7 ± 1.0 |
| Proximal radius | 0.82 ± 0.04* | 0.82 ± 0.04* | -0.6 ± 0.3* | 0.84 ± 0.02 | 0.83 ± 0.02 | -0.8 ± 0.5 |
| Total Body | 1.23 ± 0.02 | 1.22 ± 0.02 | -0.8 ± 0.3 ^a | 1.29 ± 0.04 | 1.28 ± 0.04 | -0.3 ± 0.7 |
| Pelvis | 1.25 ± 0.04 | 1.21 ± 0.04 | -3.6 ± 0.8 ^b | 1.28 ± 0.06 | 1.30 ± 0.06 | 1.6 ± 1.0 ^d |
| Legs | 1.35 ± 0.04 | 1.32 ± 0.04 | -1.9 ± 0.7 ^a | 1.48 ± 0.07 | 1.46 ± 0.06 | -0.7 ± 0.8 |
| Arms | 0.95 ± 0.02 | 0.94 ± 0.02 | -0.7 ± 0.9 | 0.91 ± 0.03 | 0.91 ± 0.02 | 0.2 ± 0.6 |

^a baseline vs. bed rest p < 0.05, ^b baseline vs. bed rest p < 0.01, ^c control vs. alendronate p < 0.05
^d control vs. alendronate p < 0.01, *5 control subjects only

Table 1. Change in bone mineral density (GM/CC) after 17 weeks of bed rest (MEAN ±SE).

| | Control (n=13) | | | Alendronate (n=8) | | |
|--------------------------------------|----------------|-------------|------------------------|-------------------|-------------|---------------------------|
| | Baseline | Bed Rest | % Change | Baseline | Bed Rest | % Change |
| NTx (nmol/mmol) | 41.0 ± 4.5 | 56.4 ± 6.0 | 40 ± 9 ^b | 36.8 ± 4.0 | 28.1 ± 3.9 | -25 ± 5 ^{b,d} |
| PYD (nmol/mmol) | 18.8 ± 1.6 | 26.7 ± 1.7 | 44 ± 4 ^b | 22.5 ± 1.6 | 28.9 ± 3.4 | 27 ± 8 ^{a,c} |
| DPD (nmol/mmol) | 5.1 ± 0.3 | 6.8 ± 0.5 | 35 ± 4 ^b | 5.5 ± 0.2 | 6.3 ± 0.3 | 16 ± 6 ^{a,c} |
| Alk Phos (U/L) | 58.3 ± 4.5 | 62.5 ± 4.9 | 8 ± 5 | 66.3 ± 4.7 | 62.6 ± 4.3 | -5 ± 1 ^{a,c} |
| BS Alk Phos (U/L) | 16.2 ± 1.2* | 15.2 ± 1.2 | -6 ± 5* | 17.4 ± 1.5 | 14.2 ± 1.1 | -18 ± 3 ^{b,c} |
| Osteocalcin (ng/ml) | 11.5 ± 1.2 | 12.9 ± 1.2 | 18 ± 11 | 13.3 ± 1.1 | 11.3 ± 0.9 | -14 ± 5 ^{a,c} |
| 1,25 (OH) ₂ Vit D (pg/ml) | 28.4 ± 2.9 | 24.5 ± 3.6 | -15 ± 6 | 40.2 ± 3.6 | 38.8 ± 3.5 | 3 ± 13 |
| 25 Vit D (pg/ml) | 20.8 ± 2.3* | 21.7 ± 2.1* | 8.4 ± 14* | 30.5 ± 3.1 | 28.5 ± 1.4 | -0.3 ± 9 |
| PTH (MM) (pg/ml) | 102.0 ± 9.4* | 89.1 ± 8.2* | -6 ± 2 | 112.0 ± 8.3 | 114.6 ± 8.8 | 4 ± 7 |
| PTH (IM) (pg/ml) | 23.5 ± 2.1* | 17.9 ± 1.4* | -23 ± 6 ^{*,a} | 24.7 ± 3.0 | 30.3 ± 2.9 | 31 ± 16 ^c |
| Serum Ca (mg/dl) | 9.6 ± 0.1 | 9.7 ± 0.1 | 1.8 ± 1.0 | 9.8 ± 0.1 | 9.3 ± 0.04 | -4.3 ± 1.1 ^{b,d} |
| Ionized Ca (mmol/L) | 1.26 ± 0.01 | 1.27 ± 0.01 | 0.5 ± 0.4 | 1.24 ± 0.01 | 1.21 ± 0.01 | -2.0 ± 0.6 ^{a,d} |

^a baseline vs. bed rest p < 0.05, ^b baseline vs. bed rest p < 0.01, ^c control vs. alendronate p < 0.05,
^d control vs. alendronate p < 0.01, * 5 control subjects only

Abbreviations: NTx, n-telopeptide in nanomoles/millimole of creatinine; PYD, pyridinium crosslinks in nanomoles/millimole of creatinine; DPD, deoxypyridinoline crosslinks in nanomoles/millimole of creatinine; Alk phos, alkaline phosphatase in units/liter; BS Alk Phos, bone specific alkaline phosphatase in units/liter; 1,25 (OH)₂ Vit D, 1,25 Dihydroxyvitamin D; 25 Vit D, 25 Hydroxyvitamin D; PTH (MM), parathyroid hormone mid-molecule; PTH (IM) parathyroid intact molecule.

Table 2. Change in bone metabolism markers and regulatory hormones during 17 weeks of bed rest (MEAN ±SE).

ionized Ca decreased slightly in the treatment group.

Table 3 gives the fecal and urine Ca excretion and balance in the treated and control groups. Compared to baseline, urinary Ca excretion increased on average 72 mg/day in controls while the treated group demonstrated a decrease of 24 mg/day. The difference in urinary Ca excretion between control and treated groups was significant. On the other hand, fecal Ca excretion showed a somewhat different response. While the controls showed a significant increase of 147 mg/day, the treated group also showed an increase of 86 mg/day although it did not reach significance. Figures 4-6 give the mean percentage change in urinary Ca, fecal Ca and Ca balance for the control and treated groups for the 3 measured time points during bed rest. Fecal Ca in the treated group appears to be increasing with length of bed rest, however the differences between time points was not significant. The Wilk's Lambda statistic showed that the between group changes of the combined urine and fecal excretion were significant ($p=0.0145$). The overall mean Ca balance was -213 mg/day in the controls and a non-significant -62 mg/day in the treatment group.

Table 4 gives the lean body mass measurements obtained from the DEXA measurements. Lean body mass was only available in 6 of the 8 previous subjects resulting in 11 controls. The lean tissue in the legs and total body showed similar decreases, although the whole body change in the alendronate group was not significant. The arms show a non-significant increase in both groups. This table demonstrates that the LBM changes in the control and treated groups were similar.

Discussion

This bed rest study demonstrated that alendronate ingestion is an effective countermeasure during bed rest, and likely in weightlessness as well. Thirteen untreated bed rest subjects show loss of bone mass in all regions with the exception of the wrist and arms. Compared to baseline, the alendronate treated subjects show no loss in BMD except the calcaneus which appears to be half that seen in the controls. All 3 bone resorption markers were elevated in the controls while decreased or elevated significantly less in the treated group. Formation markers, unchanged in the controls, are decreased in the treated group as one would expect, if alendronate is suppressing remodeling. Net Ca balance was improved by alendronate treatment. The data indicate, however, that alendronate may not completely prevent bone loss. For example, in the treated group, there was loss in the calcaneus, the pyridinoline markers were elevated, fecal Ca appears elevated and net Ca balance is negative - the latter two did not reach significance. Also this study lasted only 4 months, while missions to the planets may last many years. Extrapolating effectiveness much beyond




Figure 1. Mean \pm SE percent change in NTx for the control and treated groups for the 3 measured time points during bed rest.




Figure 2. Mean \pm SE percent change in osteocalcin for the control and treated groups for the 3 measured time points during bed rest.




Figure 3. Mean \pm SE percent change in intact molecule PTH for the control and treated groups for the 3 measured time points during bed rest.

| | Control (n=13) | | Alendronate (n = 8) | |
|-------------------------|----------------|------------------------|---------------------|-----------------------|
| | Pre-bed rest | Bed rest | Pre-bed rest | Bed rest |
| Ca intake (mg/day) | 1002 ± 12 | 1007 ± 13 | 1026 ± 9 | 1027 ± 12 |
| Urine Ca (mg/day) | 176 ± 20 | 248 ± 27 ^b | 187 ± 35 | 163 ± 29 |
| Fecal Ca (mg/day) | 724 ± 42 | 871 ± 30 ^b | 707 ± 43 | 793 ± 24 |
| Ca balance (mg/day) | 102 ± 49 | -112 ± 37 ^b | 132 ± 34 | 70 ± 30 |
| Net Ca balance (mg/day) | | -213 ± 36 ^b | | -62 ± 38 ^c |
| Delta Urine Ca (mg/day) | | 72 ± 8 ^b | | -24 ± 13 ^d |
| Delta Fecal Ca (mg/day) | | 147 ± 36 ^b | | 86 ± 37 |

^a baseline vs. bed rest p<0.05, ^b baseline vs. bed rest p<0.01, ^c control vs. alendronate p<0.05, ^d control vs. alendronate p<0.01

Table 3. Calcium excretion and balance (MEAN ±SE).

| | Control (n=11) | | | Alendronate (n = 8) | | |
|-----------------|----------------|---------------|-------------------------|---------------------|---------------|-------------------------|
| | Baseline | Post-Bed Rest | % Change | Baseline | Post-Bed Rest | % Change |
| Whole Body (kg) | 66.4 ± 3.7 | 64.3 ± 3.6 | -3.0 ± 1.1 ^a | 55.2 ± 1.9 | 54.3 ± 1.8 | -1.6 ± 0.9 |
| Legs | 19.0 ± 0.8 | 17.4 ± 0.7 | -8.5 ± 1.2 ^b | 16.9 ± 0.7 | 15.5 ± 0.7 | -8.4 ± 1.0 ^b |
| Trunk | 29.7 ± 1.4* | 29.1 ± 1.2* | -1.6 ± 1.5* | 27.5 ± 0.9 | 27.7 ± 0.7 | 0.6 ± 1.1 |
| Arms | 7.6 ± 0.4 | 7.8 ± 0.4 | 2.5 ± 1.4 | 6.8 ± 0.4 | 7.0 ± 0.4 | 3.0 ± 1.7 |

^a baseline vs. bed rest p < 0.05, ^b baseline vs. bed rest p < 0.01, * 9 control subjects only. There were no significant differences between the control and alendronate group.

Table 4. Change in lean body mass (kg) after 17 weeks of bed rest (MEAN ±SE).

the study duration is not warranted at this time. Therefore, research designed to elicit the basic mechanism(s) is recommended. Also, it should be mentioned that none of the bone parameters reported in this paper evaluate possible architectural changes within bone, an important component of bone strength. Quantitative computerized tomography (QCT) measurements of the spine, hip or lower extremities would provide valuable data in this regard.

Other than the one subject with a hiatal hernia, described in the methods, the daily ingestion of the alendronate pill was well tolerated over the 4 months in bed rest. All bisphosphonates have a theoretical potential of producing hypocalcemia by inhibiting bone resorption. In one study, bed rest subjects were given 20 mg/day of alendronate for two weeks prior to and during three weeks of bed rest. While ambulatory, two of eight subjects developed mild, asymptomatic hypocalcemia that reversed when the subjects went into bed rest¹⁴. In the current study, where the drug was given during bed rest, only a small decrease in total serum and ionized Ca, 4.2% and 2% respectively, were observed.

The DPD and PYD were elevated compared to baseline in the treated group, but significantly less compared to the control subjects. The difference in NTx response between the two groups is much more pronounced. This probably is an indication that NTx is a more specific index of bone resorption, while DPD and PYD (as measured by ELISA) are confounded, especially in cases where there is coincident muscle loss (non-bone collagen breakdown). There was also discordance between the intact and mid-molecule PTH assay results. It would appear that the intact assay better reflects the changes in total and ionized serum Ca.

From ground-based human bed rest studies it appears that decreased muscle forces result in an uncoupling of bone resorption and formation which leads to rapid bone loss^{6,8}. In these ground-based studies, resorption markers increased very early after the onset of bed rest and remained elevated for the duration of bed rest (120 days). Formation markers, on the other hand, indicate bone formation increasing slowly or not at all. During space flight, crew members exhibit a rapid and sustained increase in resorption markers that is accompanied

by continuous Ca loss, similar to bed rest^{6,20}. While formation markers during bed rest are unchanged, less published flight data is available concerning formation markers, but the limited data does not indicate an increase. In any case, there does appear to be an uncoupling similar to bed rest, the exact mechanism of which is not completely understood. Therefore, a strategy that targets the increased osteoclastic activity would seem to be a promising approach. The results of this study confirm that the uncoupling of bone remodeling is a critical part of the underlining mechanism for the loss of bone in disuse. Further, these results indicate that reducing remodeling using alendronate can prevent this loss by uncoupling the usual mechanical relationship between muscle and bone, i.e., disuse leads to muscle and bone atrophy.

For short duration missions on the order of a few weeks to a month, the amount of bone that can be lost is of little concern, especially if lost bone is eventually recovered. With increasingly lengthy flights, such as tours aboard the space station and trips to nearby planets, the amount and distribution of lost bone becomes important. The Russians, having operated the space station Mir for many years, recognized this problem and implemented a comprehensive countermeasure program. This program consisted of several hours of exercise a day to maintain physical fitness through a series of exercises consisting of bungee cord resistive exercises, bicycle ergometer, and running on a treadmill²¹. Despite the comprehensive nature of this program and its success in ameliorating many untoward effects of weightlessness, its effect on bone was obviously not complete since crew members lost substantial bone during flight²⁻³. However, it is unknown what the loss would have been without this countermeasure program since a non-exercising control group was not possible. Also, there were individual differences in crew compliance performing the exercise regime as a result of personal preference or mission constraints²². For example, exercise is generally not practical during the first days of flight because of scheduling constraints early in the mission, while many crew members increased their exercise intensity in the weeks just prior to return to earth. Reports from returning crew also indicate significant variability among crew members in the amount, scheduling and type of exercise performed during flight. Without disregarding the possibility that the exercise program may have prevented some bone loss, it does appear to require modification. The current exercise program carried out on the International Space Station (ISS) has modified the Russian program to include greater amounts of resistive exercise using a device that is intended to provide mechanical forces on the bones and muscles that mimic weightlifting in one G. The effectiveness of this program is currently being evaluated with ISS crew members; this regime will require many flights to determine efficacy. Even if the current exercise program is successful, which it may not be, having a safe pharmacological countermeasure available will




Figure 4. Mean \pm SE mg/day excretion of urinary Ca for the control and treated groups for the 3 measured time points during bed rest.




Figure 5. Mean \pm SE mg/day excretion of fecal Ca for the control and treated groups for the 3 measured time points during bed rest.




Figure 6. Mean \pm SE Ca balance in mg/day for the control and treated groups for the 3 measured time points during bed rest.

be important. The current exercise requires a considerable amount of time and in some situations, such as occurred aboard MIR in the late 1990's, there may not be adequate time for exercise. A combination of alendronate with exercise might reduce the time needed to protect the skeleton, but needs investigation. There may be other situations when the crew members cannot exercise, e.g., injury, equipment failure or sickness. Other forms of alendronate, e.g., the 8 day pill marketed by Merck would simplify the use of this drug in space. Newer bisphosphonates being investigated in various laboratories may prove even more effective, e.g., injectable forms that suppress remodeling for several months are appealing if shown to be safe and effective.

Acknowledgments

This work was supported by NASA grant number NCC9-61, the GCRC at Baylor College of Medicine, Houston, Texas, USA, NIH grant number M01RR00188 and the National Space Biomedical Research Institute, Baylor College of Medicine. We would also like to thank Dr. Edward Powers for the medical care provided to the subjects.

References

- Nicogossian AE, Huntoon CL, Pool SL. Space Physiology and Medicine, 3rd edition. Lea & Febiger, Philadelphia; 1994.
- Oganov VS, Grigoriev AI, Voronin LI, Rakmanov AS, Bakulin AV, Schneider V, LeBlanc A. Bone mineral density in cosmonauts after 4.5-6 month long flights aboard orbital station MIR. *Aerospace and Environmental Medicine* 1992; 26(No.5-6):20-24.
- LeBlanc A, Schneider V, Shackelford L, West S, Oganov V, Bakulin A, Voronin L. Bone mineral and lean tissue loss after long duration space flight. *J Musculoskel Neuron Interact* 2000; 1:157-160.
- LeBlanc AD, Lin C, Shackelford L, Sinityn V, Evans H, Belichenko O, Schenkman B, Kozlovskaya I, Oganov V, Bakulin A, Hedrick T, Feeback D. Muscle volume, MRI relaxation times (T2) and body composition after space flight. *J Appl Physiol* 2000; 89:2158-2164.
- Oganov V, Bakulin A, Novikov V, Murashko L, Kabitskaya O, Schneider V, Shackelford L, Evans H, LeBlanc A. Results of human bone densitometry after prolonged space flights. International Cooperation in space Station MIR. Lion, France, March 19-21, 2001 (Abstract)
- Smith SM, Nillen JL, LeBlanc AD, Lipton A, Demers LM, Lane HW, Leach CS. Collagen cross-link excretion during space flight and bed rest. *J Clin Endocrinol Metab* 1998; 83:3584-3591.
- LeBlanc A, Schneider V, Evans H, Engelbretson D, Krebs J. Bone mineral loss and recovery after 17 weeks of bed rest. *J Bone Miner Res* 1990; 5:843-850.
- LeBlanc A, Schneider V, Spector E, Evans H, Rowe R, Lane H, Demers L, Lipton A. Calcium absorption, endogenous secretion and endocrine changes during and after long-term bed rest. *Bone* 1995; 16:301S-304S.
- Lockwood DR, Vogel JM, Schneider VS, Hulley SB. Effect of the disphosphonate EHDP on bone mineral metabolism during prolonged bed rest. *J Clin Endocrinol Metab* 1975; 41:533-541.
- Schneider VS, McDonald J. Skeletal calcium homeostasis and countermeasures to prevent disuse osteoporosis. *Calcif Tissue Int* 1984; 36:S151-154.
- Fleisch H, Russell RGG, Simpson B, Muhlbauer RC. Prevention by a diphosphonate of immobilization "osteoporosis" in rats. *Nature* 1969; 223:211-212.
- Meunier PG, Chapuy MC, Delmas P. Intravenous disodium etidronate therapy in Paget's disease of bone and hypercalcemia of malignancy. *Am J Med* 1987; 82(Suppl 2A):71-78.
- Schneider VS, personal communication.
- Ruml LA, Dubois SK, Roberts ML, Pak CYC. Prevention of hypercalciuria and stone-forming propensity during prolonged bedrest by alendronate. *J Bone Miner Res* 1995; 10:655-662.
- Schneider VS. Modification of negative calcium balance and bone loss during prolonged bed rest. NASA Terminal Report No. T-660-1981, Vol 1-3; 1981.
- LeBlanc A, Evans H, Marsh C, Schneider V, Johnson P, Jhingran S. Precision of dual photon absorptiometry measurements. *J Nucl Med* 1986; 27(8):1362-1365.
- LeBlanc A, Schneider V, Engelbretson D, Evans H. Precision of regional bone mineral measurements obtained from total-body scans. *J Nucl Med* 1990; 31:43-45.
- Spector E, LeBlanc A, Shackelford L. Hologic QDR 2000 whole body scans: A comparison of three combinations of scan modes and analysis software. *Osteoporos Int* 1995; 5:440-445.
- Anderson TW. An Introduction to multivariate statistical analysis, 2nd edition, Wiley, New York; 1984.
- Rambaut PC, Johnston RS. Prolonged weightlessness and calcium loss in man. *Acta Astronautica* 1979; 6:1113-1122.
- Nicogossian AE, Sawin CF, Grigoriev AI. Countermeasures to space deconditioning. In: Nicogossian AE, Huntoon CL, Pool SL (eds). *Space Physiology and Medicine*, 3rd edition, Lea & Febiger, Philadelphia; 1994: 447-467.
- Kozlovskaya IB, Barmin VA, Stepanov VI, Kharitonov NM. Results of studies of motor functions in long-term space flights. *Physiologist* 1990; 33(1): S1-S3.